

Complete Summary

GUIDELINE TITLE

Unstable angina pectoris.

BIBLIOGRAPHIC SOURCE(S)

Finnish Medical Society Duodecim. Unstable angina pectoris. In: EBM Guidelines. Evidence-Based Medicine [CD-ROM]. Helsinki, Finland: Duodecim Medical Publications Ltd.; 2006 Mar 15 [Various].

GUIDELINE STATUS

This is the current release of the guideline.

This guideline updates a previous version: Finnish Medical Society Duodecim. Unstable angina pectoris. In: EBM Guidelines. Evidence-Based Medicine [CD-ROM]. Helsinki, Finland: Duodecim Medical Publications Ltd.; 2004 Sep 14 [Various].

COMPLETE SUMMARY CONTENT

SCOPE
 METHODOLOGY - including Rating Scheme and Cost Analysis
 RECOMMENDATIONS
 EVIDENCE SUPPORTING THE RECOMMENDATIONS
 BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS
 IMPLEMENTATION OF THE GUIDELINE
 INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT
 CATEGORIES
 IDENTIFYING INFORMATION AND AVAILABILITY
 DISCLAIMER

SCOPE

DISEASE/CONDITION(S)

Unstable angina pectoris

GUIDELINE CATEGORY

Diagnosis
 Evaluation
 Risk Assessment
 Treatment

CLINICAL SPECIALTY

Cardiology
Family Practice
Internal Medicine

INTENDED USERS

Health Care Providers
Physicians

GUIDELINE OBJECTIVE(S)

Evidence-Based Medicine Guidelines collects, summarizes, and updates the core clinical knowledge essential in general practice. The guidelines also describe the scientific evidence underlying the given recommendations.

TARGET POPULATION

Patients with suspected or confirmed unstable angina

INTERVENTIONS AND PRACTICES CONSIDERED

Diagnosis

1. Electrocardiogram
2. Assessment of signs and symptoms
3. Measurement of myocardial markers, such as troponin T or troponin I and creatine kinase isoenzyme MB mass (CK-MBm)

Treatment

1. Aspirin
2. Clopidogrel in addition to aspirin
3. Nitrate
4. Beta-blocker (metoprolol)
5. Low-molecular-weight (LMW) heparins (such as dalteparin) simultaneously with aspirin
6. Glycoprotein IIb/IIIa inhibitors in selected patients
7. Cardiac monitoring
8. Angiography
9. Revascularization (clopidogrel and intravenous glycoprotein IIb/IIIa (GP IIb/IIIa) inhibitor in addition to aspirin and low-molecular-weight heparin while waiting for the procedure)
10. Thrombolytic therapy
11. Percutaneous transluminal angioplasty (PTA) with insertion of a stent followed by clopidogrel in combination with aspirin
12. Elimination of risk factors

Evaluation

1. Electrocardiogram
2. Measurement of myocardial markers, such as troponin T or troponin I and creatine kinase isoenzyme MB mass
3. Exercise testing

MAJOR OUTCOMES CONSIDERED

- Mortality
- Incidence of myocardial infarction
- Efficacy of treatment for angina in reducing mortality and incidence of myocardial infarction
- Episodes of chest pain and severity of chest pain (i.e., need for sublingual nitroglycerin)
- Incidence of ischaemic events/recurrent angina
- Bleeding complications
- Need for urgent revascularization
- Sensitivity and specificity of troponin I and T for predicting adverse cardiac events in unstable angina pectoris

METHODOLOGY

METHODS USED TO COLLECT/SELECT EVIDENCE

Hand-searches of Published Literature (Primary Sources)
 Hand-searches of Published Literature (Secondary Sources)
 Searches of Electronic Databases

DESCRIPTION OF METHODS USED TO COLLECT/SELECT THE EVIDENCE

The evidence reviewed was collected from the Cochrane database of systematic reviews and the database of abstracts of reviews of effectiveness (DARE). In addition, the Cochrane Library and medical journals were searched specifically for original publications.

NUMBER OF SOURCE DOCUMENTS

Not stated

METHODS USED TO ASSESS THE QUALITY AND STRENGTH OF THE EVIDENCE

Weighting According to a Rating Scheme (Scheme Given)

RATING SCHEME FOR THE STRENGTH OF THE EVIDENCE

Levels of Evidence

- A. Strong research-based evidence. Multiple relevant, high-quality scientific studies with homogenic results.

- B. Moderate research-based evidence. At least one relevant, high-quality study or multiple adequate studies.
- C. Limited research-based evidence. At least one adequate scientific study.
- D. No research-based evidence. Expert panel evaluation of other information.

METHODS USED TO ANALYZE THE EVIDENCE

Review of Published Meta-Analyses
Systematic Review

DESCRIPTION OF THE METHODS USED TO ANALYZE THE EVIDENCE

Not stated

METHODS USED TO FORMULATE THE RECOMMENDATIONS

Not stated

RATING SCHEME FOR THE STRENGTH OF THE RECOMMENDATIONS

Not applicable

COST ANALYSIS

A formal cost analysis was not performed and published cost analyses were not reviewed.

METHOD OF GUIDELINE VALIDATION

Peer Review

DESCRIPTION OF METHOD OF GUIDELINE VALIDATION

Not stated

RECOMMENDATIONS

MAJOR RECOMMENDATIONS

The levels of evidence [A-D] supporting the recommendations are defined at the end of the "Major Recommendations" field.

Objective

- To recognize ischaemic chest pain that may be prodromal to acute infarction (acute coronary syndrome [ACS]) and to accompany the patient to a cardiac monitoring unit for active drug treatment or rapid revascularization (Wallentin et al., 2000) [B].

- To assess the patient's risk factors for coronary heart disease, and to refer high risk patients readily to a hospital for coronary angiography. The only symptom of ischaemia may be a general deterioration in patient's physical condition, or dyspnoea, without chest pain.

Definition

- Symptom complexes caused by a sudden narrowing or obstruction of a coronary artery are called acute coronary syndromes (ACS). These include
 - Unstable angina pectoris
 - Myocardial infarction without ST-elevation
 - Myocardial infarction with ST-elevation (See the Finnish Medical Society Duodecim guideline "Thrombolytic therapy and balloon angioplasty in acute ST elevation myocardial infarction [STEMI]")
- The most common cause of an acute coronary syndrome is a tear in an atheroma plaque and a thrombus that is formed on it.

Risk Groups and Clinical Signs

- The presence of a marker (cardiac troponin T and I, creatine kinase isoenzyme MB mass [CK-MBm]) is the single most important predictor for future coronary events.
 - Marker-positive patients are referred to angiography and revascularization.
 - Marker-negatives are referred to exercise tolerance test.
- Unstable angina pectoris (UAP) is a heterogeneous group of diseases covering the range between stable angina pectoris (AP) and acute myocardial infarction (AMI).
- New (sudden) AP in a high-risk patient is always a serious condition.
- An aggravation in stable AP to unstable AP always necessitates a reassessment of risk and often a change in the line of treatment.
- There may not always be pain; rather, the main symptom is a decrease in exercise tolerance (sudden decrease in physical fitness) or acute left ventricle failure.
- In the electrocardiogram (ECG) an ST segment depression precedes the pain. Symptomless (silent) ischaemia in a patient at risk is a significant finding. Ischaemia may not always be visible in a resting ECG. An ECG registered while the patient has pain is invariably valuable.
- The border between UAP and T-wave infarction (non-Q infarction) is shifting. For example, very proximal occlusion in the left anterior descending artery (LAD) causes a symmetric T inversion in chest leads. Elevation of myocardial markers indicates that the patient has an infarction.

Treatment

- Treatment is normally carried out in a cardiac monitoring unit.
- Pharmacological treatment should be started in the first point of care.
- The mildest form (recent angina) can be treated in a health care centre ward under careful monitoring. Remember the risk of myocardial infarction (MI). The risk diminishes with time as the angina stabilizes.

Anti-ischaemic and Antithrombotic Treatment

- All patients with suspected unstable angina (no changes in ECG or myocardial markers)
 - Aspirin 100 milligrams (mg) per day continuously, unless there are contraindications (Natarajan, 2002; "Collaborative overview," 1994) [A].
 - Clopidogrel together with aspirin: first a loading dose of 300 mg, then 75 mg per day.
 - Nitrate (Natarajan, 2000) [D] intravenously (See the Finnish Medical Society Duodecim guideline "Nitrate infusion in angina pectoris and myocardial infarction") or orally according to the situation.
 - Beta-blocker (Natarajan, 2002) [C] (metoprolol). Heart rate should be 50 to 60 beats per minute and systolic pressure below 150 mmHg.
 - Low-molecular-weight (LMW) heparin (Zed, Tisdale, & Borzak, 1999; Nicholson, Milne, & Stein, 2000; The Health Technology Assessment Database, HTA-20000891, 2001) [A] (e.g., dalteparin 100-120 IU/kg x 2 daily for one week) is given simultaneously with aspirin. The treatment can be continued with half the dose for about 1 month. Unstable AP patients with an elevated troponin T concentration derive the greatest benefit from the treatment (LMW heparin + aspirin). Pharmacotherapy and invasive treatment do not exclude one another.
 - Glycoprotein IIb/IIIa inhibitors for selected high risk patient groups in hospital care
 - There is no benefit from fibrinolytic therapy.
- High-risk patients: Unstable angina and ischaemia on ECG or elevated myocardial markers (Olatidoye et al., 1998) [A], acute left ventricle failure (lung oedema, mitral regurgitation, hypotension)
 - Immediate angiography and revascularization. While waiting for the procedure, the thrombosis should be stabilized with clopidogrel (an initial dose of 300 mg before transportation, thereafter 75 mg daily) and an intravenous (i.v.) glycoprotein IIb/IIIa (GP IIb/IIIa) inhibitor (Natarajan, 2002) [B] in addition to aspirin and LMW heparin . (Fibrinolytic treatment has no effect on a vessel obstruction caused by aggregated platelets.)
- Thrombolytic therapy or immediate percutaneous transluminal angioplasty (PTA) (during which a stent can be inserted) is indicated if ECG reveals a transmural injury. (See article on revascularization: Evidence-Based Medicine guideline, "Coronary heart disease symptoms, diagnosis and treatment.") After the insertion of a stent, clopidogrel is used in combination with aspirin for one month.
- Further treatment of patients with symptoms or signs of ischaemia on ECG and normal myocardial markers
 - Symptom-limited exercise test performed within 2 to 4 days.
 - If the patient has symptoms or signs of ischaemia during the exercise test or signs in ECG at a low pulse-pressure product, refer immediately to angiography.
 - In case of no symptoms or signs of ischaemia during light exercise or no signs in ECG, or if they occur only with a high pulse-pressure product, begin conservative treatment and elimination of risk factors. Prophylaxis can be intensified by adding clopidogrel to aspirin.

Organizing Treatment

- UAP is a serious but often curable syndrome. A well-organized care pathway and efficient follow-up care ensures that the appropriate treatment can be given rapidly.

Related Evidence

- Intravenous heparin combined with aspirin is probably effective compared to aspirin alone in reducing myocardial infarction or death, but the number of patients studied is too small for statistical significance (Oler et al., 1996) [C].

Definitions:

Levels of Evidence

- A. Strong research-based evidence. Multiple relevant, high-quality scientific studies with homogenic results.
- B. Moderate research-based evidence. At least one relevant, high-quality study or multiple adequate studies.
- C. Limited research-based evidence. At least one adequate scientific study.
- D. No research-based evidence. Expert panel evaluation of other information.

CLINICAL ALGORITHM(S)

None provided

EVIDENCE SUPPORTING THE RECOMMENDATIONS

REFERENCES SUPPORTING THE RECOMMENDATIONS

[References open in a new window](#)

TYPE OF EVIDENCE SUPPORTING THE RECOMMENDATIONS

Concise summaries of scientific evidence attached to the individual guidelines are the unique feature of the Evidence-Based Medicine Guidelines. The evidence summaries allow the clinician to judge how well-founded the treatment recommendations are. The type of supporting evidence is identified and graded for select recommendations (see the "Major Recommendations" field).

BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS

POTENTIAL BENEFITS

- Appropriate diagnosis and treatment of unstable angina pectoris
- Appropriate assessment of coronary heart disease risk factors
- Reduced rate of myocardial infarction and death

POTENTIAL HARMS

Low-molecular-weight heparin can cause bleeding complications.

IMPLEMENTATION OF THE GUIDELINE

DESCRIPTION OF IMPLEMENTATION STRATEGY

An implementation strategy was not provided.

INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT CATEGORIES

IOM CARE NEED

Getting Better
Staying Healthy

IOM DOMAIN

Effectiveness
Timeliness

IDENTIFYING INFORMATION AND AVAILABILITY

BIBLIOGRAPHIC SOURCE(S)

Finnish Medical Society Duodecim. Unstable angina pectoris. In: EBM Guidelines. Evidence-Based Medicine [CD-ROM]. Helsinki, Finland: Duodecim Medical Publications Ltd.; 2006 Mar 15 [Various].

ADAPTATION

Not applicable: The guideline was not adapted from another source.

DATE RELEASED

2001 Apr 30 (revised 2006 Mar 15)

GUIDELINE DEVELOPER(S)

Finnish Medical Society Duodecim - Professional Association

SOURCE(S) OF FUNDING

Finnish Medical Society Duodecim

GUIDELINE COMMITTEE

Editorial Team of EBM Guidelines

COMPOSITION OF GROUP THAT AUTHORED THE GUIDELINE

Primary Authors: Editors

FINANCIAL DISCLOSURES/CONFLICTS OF INTEREST

Not stated

GUIDELINE STATUS

This is the current release of the guideline.

This guideline updates a previous version: Finnish Medical Society Duodecim. Unstable angina pectoris. In: EBM Guidelines. Evidence-Based Medicine [CD-ROM]. Helsinki, Finland: Duodecim Medical Publications Ltd.; 2004 Sep 14 [Various].

GUIDELINE AVAILABILITY

This guideline is included in a CD-ROM titled "EBM Guidelines. Evidence-Based Medicine" available from Duodecim Medical Publications, Ltd, PO Box 713, 00101 Helsinki, Finland; e-mail: info@ebm-guidelines.com; Web site: www.ebm-guidelines.com.

AVAILABILITY OF COMPANION DOCUMENTS

None available

PATIENT RESOURCES

None available

NGC STATUS

This summary was completed by ECRI on August 28, 2001. The information was verified by the guideline developer as of October 26, 2001. This summary was updated by ECRI on December 9, 2002, April 2, 2004, February 22, 2005, and May 25, 2006.

COPYRIGHT STATEMENT

This NGC summary is based on the original guideline, which is subject to the guideline developer's copyright restrictions.

DISCLAIMER

NGC DISCLAIMER

The National Guideline Clearinghouse™ (NGC) does not develop, produce, approve, or endorse the guidelines represented on this site.

All guidelines summarized by NGC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public or private organizations, other government agencies, health care organizations or plans, and similar entities.

Guidelines represented on the NGC Web site are submitted by guideline developers, and are screened solely to determine that they meet the NGC Inclusion Criteria which may be found at <http://www.guideline.gov/about/inclusion.aspx>.

NGC, AHRQ, and its contractor ECRI make no warranties concerning the content or clinical efficacy or effectiveness of the clinical practice guidelines and related materials represented on this site. Moreover, the views and opinions of developers or authors of guidelines represented on this site do not necessarily state or reflect those of NGC, AHRQ, or its contractor ECRI, and inclusion or hosting of guidelines in NGC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding guideline content are directed to contact the guideline developer.

© 1998-2006 National Guideline Clearinghouse

Date Modified: 9/25/2006

